

Scoring Categories

- **Category 1:** Patient may be controlled on the current treatment regimen. Continue monitoring the patient based on best medical treatment and your professional judgment/clinical guidelines.
- **Category 2:** Patient may not be controlled on the current treatment regimen. However, additional benefits may be obtained from further treatment optimisation. Device-aided therapies may not be needed at this time. However, use your patient's medical history, treatment preference and your best medical judgment for treatment recommendation.
- **Category 3:** Patient may not be controlled on the current treatment regimen and may benefit from device-aided therapy. It is suggested that you evaluate eligibility for device-aided therapy based on patient's medical history, treatment preference and your best medical judgment for treatment recommendation.

Section 1 Scoring

Screeners for inadequate control on current orals

- Taking oral levodopa ≥ 5 times/day
- ≥ 2 hours of “off” time/day
- Presence of unpredictable fluctuations of motor symptoms
- Presence of troublesome dyskinesia
- Limitations in ≥ 1 Activities of Daily Living

If answer is NO to ALL the questions in Section 1 then patient is classified as [Category 1](#).

If answer is YES to ANY questions in Section 1 then patient continues to Section 2.

Section 2 Scoring

Frequency score

- 0 Never
- 1 Rarely
- 2 Sometimes
- 3 Most/all the time

Severity score

- 1 Mild
- 2 Moderate
- 3 Severe

Level of independence (Activities of Daily Living)

- 0 Independent in all activities
- 1 Independent in most activities
- 2 Needs assistance/dependent in some activities
- 3 Totally dependent in all activities

Falls per month

- 0 No Falls
- 1 1
- 2 2 or more

- If in **Section 1**, a levodopa dose ≤ 3 was selected, **AND** “Yes” was selected for **ANY** of the other categories (eg, ≥ 2 hours “Off” time/day, presence of unpredictable motor fluctuations, presence of troublesome dyskinesia, limitations in activities of daily living) the patient will be classified as [Category 2](#), regardless of frequency/severity of symptoms.
- If “Yes” was selected for **ANY** of the categories (eg, ≥ 2 hours “Off” time/day, presence of unpredictable motor fluctuations, presence of troublesome dyskinesia, limitations in activities of daily living) in **Section 1**, **AND** a levodopa dose ≥ 4 was selected, please refer to the algorithm on the following page for **Category Classification**

≥4 doses of levodopa and:

(Frequency x Severity) score of ≥ 3 for:

- Unpredictable motor fluctuations*
- Non-motor “Off” symptoms
- Freezing of Gait during “Off” time
- Hallucinations/Psychosis without insight

Category

3

(Frequency x Severity) score of < 3 for:

- Unpredictable motor fluctuations*
- Non-motor “Off” symptoms
- Freezing of Gait during “Off” time
- Hallucinations/Psychosis without insight

Category

2

A score of 2 or 3 for:

- “Off” time periods*
- Troublesome dyskinesia*
- Activities of Daily Living impairment*

Category

3

*These questions will be skipped in Section 2 if the answer in Section 1 is No.

≥4 doses of levodopa and:

A score of < 2 for:

- “Off” time periods*
- Troublesome dyskinesia*
- Activities of Daily Living impairment*

Category

2

A score of 1 or 2 for:

- Falls

Category

3

A score of 0 for:

- Falls

Category

2

*These questions will be skipped in Section 2 if the answer in Section 1 is No.

- **Please note:** The scores for “Dystonia with Pain” and “Impulse Control Disorder” have been provided for your information so that you can consider these domains when evaluating your patient for treatment recommendations. The severity and frequency of these scores do not affect the outcome of the instrument.